




Smoking status at time of delivery

% of mothers who are smokers at the time of delivery (2017/18)



Teenage mothers

% of delivery episodes where the mother is aged under 18 years in 2016/17



BME Deliveries

% of deliveries to mothers from Black and Minority Ethnic (BME) groups in 2016/17



Low birth weight of term babies

% of all live births at term in 2016 with low birth weight (<2500g)




55




Dtap/ IPV/ Hib coverage (1 year)

% of eligible children who received 3 doses of Dtap/ IPV/ Hib vaccination by their 1st birthday (2017/18)




Respiratory tract infection admissions aged 1 year

Rate of emergency admissions in infants per 10,000 population (2016/17)




Infant Mortality

Rate of deaths in infants aged under 1 year per 1,000 live births (2015 - 17)




Breastfeeding prevalence

% of all infants due a 6-8 week check that are totally or partially breastfed (2017/18)




MMR – 1st dose

% of children who received one dose of MMR vaccine on or after their 1st birthday and anytime to their 2nd birthday (2017/18)



Respiratory tract infection admissions 2,3 and 4 years

Rate of emergency admissions in infants per 10,000 population (2016/17)



MMR – 2nd dose

% of children who received two doses of MMR vaccine on or after their 1st birthday and at anytime to 5th birthday (2017/18)



The latest available data is shown at a local authority level and compared against the England figure and the comparator group, least deprived decile (IMD 2015) figures:

- Significantly better
- Significantly worse
- Significantly lower
- No significant difference
- Significantly higher
- Not comparable/Value unknown

Where a national target has been set, data has been compared against this target.

Source:
[Public Health England Fingertips Profiles](#)



School readiness

% of children achieving a good level of development at the end of Year R (2016-17)



Free school meals

% of pupils eligible for and claiming free school meals who attend state funded school (2018)




Obesity and overweight (Year R)

% of children in Year R who are obese and overweight (2017/18)



Pupils with special education needs (SEN)

% of school pupils with special education needs (2018)




Sedentary for 7 hours or more per day

% with a mean daily sedentary time in the last week over 7 hours per day (2014-15)



Physically Active (at least 1 hour every day in last week)

% physically active for at least one hour per day seven days a week (2014-15)




Obesity and overweight (Year 6)

% of children in Year 6 who are obese and overweight (2017/18)




First time entrants to Youth Justice System (10-17 year olds)

Rate per 100,000 population (2017)




Regular drinker of alcohol (every week)

% of 15 year old who have an alcoholic drink 'at least once a week' Way survey (2014-15)




Smoking prevalence (current smokers)

% of smoking prevalence at age 15 Way survey (2014-15)



Drug Use (in the last month)

% of 15 years old 'who took drugs' in the last month' Way survey (2014-15)



The latest available data is shown at a local authority level. This is compared against the England figure and the comparator group, least deprived decile (IMD 2015):

- Significantly better
- Significantly worse
- No significant difference
- Not comparable/Value unknown

Where a national target has been set, data has been compared against this target.

Source:


[Public Health Fingertips Profiles](#)



GCSEs achieved (5A*-C including English & Maths)

Educational attainment (5 or more GCSEs): % of all children (2015-16)

Wokingham:	● 70.9%
Comparator:	● 63.5%
England:	● 57.8%



Under 18 s conceptions

Rate of conceptions for females aged 15-17 per 1,000 population (2016)

Wokingham:	↓ 8.1
Comparator:	● 11.9
England:	● 18.8



Admissions as a result of self-harm (15 to 19 year olds)

Rate of admissions as a result of self-harm per 100,000 population (2016/17)

Wokingham:	↑ 720.6
Comparator:	● 652.3
England:	● 617.1



Chlamydia screened (15-24 year olds)

Proportion of population screened for chlamydia in specialist and non-specialist sexual health services (2017)

Wokingham:	↕ 15.0
Comparator:	● 16.4
England:	● 19.3

57



Under 18s conceptions leading to abortion

% of conception aged under 18 years that led to an abortion (2016)

Wokingham:	↔ 56.5%
Comparator:	● 61.6%
England:	● 51.8%



NEET (16 to 17 year olds)

% of 16-17 year olds who are NEET or whose activity is not known (2016)

Wokingham:	● 4.4%
Comparator:	● 5.5%
England:	● 6.0%

The latest available data is shown at a local authority level. This is compared against the England figure and the comparator group, least deprived decile (IMD 2015):

- Significantly better than England
- Significantly worse than England
- No significant difference to England
- Not comparable/Value unknown


Where a national target has been set, data has been compared against this target.

Source:

[Public Health Fingertips Profiles](#)

**Smoking Prevalence (current smokers)**


% of smoking prevalence among persons 18+ years from the APS survey (2017)

**Adult drinkers of alcohol (18+ years)**

% of adults drinking over 14 units of alcohol a week (2011-14)

**Physically active and inactive (19+)**

% of adults who are physically active / inactive by CMO recommendations (2016/17)

**Obesity and overweight**

% of adults (aged 18+) classified as overweight or obese (2016/17)

**Preventable Mortality (all ages)**

Rate of mortality from causes considered preventable per 100,000 population (2015/17)

**Diabetes**

Estimated diagnosis rate for people with diabetes aged 17 and over (2017)

**Suicide 10 years+**

Age-standardised mortality rate from suicide and injury of undetermined intent per 100,000 population (2015/17)

**Depression or Anxiety 18+ years**


% reporting depression or anxiety completing GP Patient Survey (2016/17)

**Bowel cancer screening (60-74 years)**

% of people eligible for bowel screening who were screened (2017)

**Cervical Screening**

% of female, 25-64, attending cervical screening within target period (3.5 or 5.5 year coverage) 2017/18

**Breast cancer screening (53-70 years)**

% of eligible women screened adequately with the previous 3 years on 31st March (2017)



The latest available data is shown at a local authority level and compared against the England figure and the comparator group, least deprived decile (IMD 2015) figures:

- Significantly better
- Significantly worse
- No significant difference
- Significantly higher
- Not comparable/Value unknown

Where a national target has been set, data has been compared against this target.


Sources:

[Public Health England Fingertips Profiles](#)
[Cervical Screening Indicator](#)





Health related quality of life

Average health status score for adults age 65 and over (2016/17)



Life expectancy at 65

Average number of years a person would expect to live (2015/17)




Excess winter deaths

Measured as the ratio of extra deaths from all causes that occur in the winter months (Aug 2014 - Jul 2017) (3 years, All Ages)



Flu vaccine

% of eligible adults aged 65+ who have received the flu vaccine (2017/18)



59



Preventable sight loss (persons 65+)

Rate of sight loss due to age related macular degeneration (AMD) in people aged 65+ per 100,000 population (2016/17)



Dementia: Recorded prevalence

Number of people with dementia recorded as a proportion of people aged 65+ registered at each GP Practice (2017)




Delayed transfer of care

Rate of delayed transfer of care 18+ per 100,000 population (2017/18)



Permanent admissions to residential and nursing homes

Rate per 100,000 population aged 65+ (2013/14)



Hospital admissions due to fall

Rate of emergency hospital admissions for injuries due to falls in people aged 65+ per 100,000 population (2017/18)



Hip fractures (65+)

Rate of emergency admissions for fractured neck of femur per 100,000 population (2017/18)



Cardiovascular Disease

Rate of deaths from cardiovascular disease among people aged 65 and over (2015/17)



The latest available data is shown at a local authority level. This is compared against the England figure and the comparator group, least deprived decile (IMD 2015):

- Significantly better
- Significantly worse
- No significant difference
- Not comparable/Value unknown

Where a national target has been set, data has been compared against this target.

Sources:
[Public Health England Fingertips Profiles](#)
[Delayed transfers of care](#)

This page is intentionally left blank